

microvilli and tonofilaments. The derivation of all lung cancer from a common endodermic stem cell is responsible for the frequent multidirectional differentiation in lung tumors, including neuroendocrine phenotype in NSCLC and squamous or glandular signs of differentiation in authenticated neuroendocrine tumors. Only three cell types are candidates for cell division and subsequent proliferation following injury: supra basal intermediate bronchial cells with small mucus granules, neuroendocrine cells and a type-II Clara cell. All three candidates themselves derive from a common endodermic totipotent stem cell which may select one or other differentiation pathway according to the set of genetic alterations which have led to its malignant transformation. The discovery that neuroendocrine cells in normal bronchial mucosa had acquired their APUD phenotype from endodermic stem cells, instead of being derived from the neural crest, helps to explain why several differentiation pathways are potentially available during clonal expansion and are simultaneously displayed in a subgroup of lung tumors.

## References

1. Travis WO, Colby TV, Corrin B at al. *Histological typing of lung and pleural tumours*. 3rd ed., World Health Organization, Geneva 1999.
2. Travis WD, Rush W, Flieder DB at al. *Survival analysis of 200 pulmonary neuroendocrine tumors with clarification of Criteria for atypical carcinoid and its separation from typical carcinoid*. Am J Surg Pathol 1998; 22(8): 934-944.
3. Roggli vL, Vollmer RT, Greenberg SD at al. *Lung cancer heterogeneity: A blinded and randomized study of 100 consecutive cases*. Human Pathol 1985; 16: 569-579.
4. Whitehead R, Cosgrove C. *Mucins and carcinoid tumors*. Pathology 1979; 11: 473-478.
5. Gould yE, Memoli VA, Dardi LE. *Multidirectional differentiation in human epithelial cancers*. J Submicrosc Cytol 1981; 13: 97-115.
6. Bolen JW, Thorning O. *Histogenetic classification of lung carcinomas. Small cell carcinomas studied by light and electron, microscopy* J Submicrosc Cytol 1982; 14: 499-514.
7. McDowell EM, Trump BF. *Pulmonary small cell carcinoma showing tripartite differentiation in individual cells*. Human Pathol 1981; 12: 286-294.
8. Hammond ME, Sause NT. *Large cell neuroendocrine tumors of the lung. Clinical significance and histopathologic definition*. Cancer 1985; 58:1624-1629.
9. Blobel GA, Gould yE, Moll R at al. *Coexpression of neuroendocrine markers and epithelial cytoskeletal proteins in bronchopulmonary neuroendocrine neoplasms*. Lab Invest 1985; 52: 39-51.
10. Brambilla E, Moro D, Gazzeri S et al. *Chemotherapy induces cell differentiation in small cell lung carcinoma*. J Clin Oncol 1991; 9: 50-61.
11. Brambilla E, Veale O, Moro D at al. *Neuro-endocrine phenotype in lung cancers: Comparison of immunocytochemistry with biochemical determination of enzyme isoenzymes*. Am J Clin Pathol 1992; 98: 88-97.
12. Brambilla E, Jacrot M, Batandier C at al. *Heterotransplantation of small cell lung carcinoma into Nude mice: Stability of phenotypic characters*. Cancer 1989; 64: 42-71.
13. Brambilla E, Brambilla C. *Heterotransplantation of small cell lung carcinoma into Nude mice: Stability of phenotypic characters*. Cancer 1989; 64: 42-71.
14. Brambilla E, Brambilla C. *Heterotransplantation of small cell lung carcinoma into Nude mice: Stability of phenotypic characters*. Cancer 1989; 64: 42-71.

1/27 ~-PA

1/2

## Divergent differentiation in thyroid tumors

M. Sobrinha-Simões<sup>1</sup>, A.R. Soares<sup>1</sup>, A.S. Rocha<sup>1</sup>,  
E. Fonseca<sup>1</sup>, M. Papotti<sup>2</sup>, R. Komminoth<sup>3</sup> and G. Bussolati<sup>1</sup>

<sup>1</sup>Institute of Molecular Pathology and Immunology University of Porto, Portugal, <sup>2</sup>Dept. of Biomedical Sciences and Human Oncology University of Turin, Italy and <sup>3</sup>University Institute of Pathology Zurich, Switzerland.

Whenever one thinks of divergent differentiation in thyroid tumors, it is the so-called mixed medullary and follicular (or papillary) carcinoma that comes to mind. The reason for this stems from the well-known dichotomy between follicular and parafollicular cells or, in other words, from the existence in the thyroid of endocrine, thyroglobulin-positive tumors and neuroendocrine, calcitonin-positive tumors. The latter part of this short review will therefore be devoted to the discussion of the diagnosis and histogenesis of mixed endocrine-neuroendocrine carcinomas of the thyroid. In the first part we will 'swim against the stream' and discuss the divergent differentiation of tumors belonging to the group of thyroglobulin-positive thyroid carcinomas. Somewhat provocatively we will focus, respectively, on the follicular variant of papillary carcinoma, considered an example of a peculiar morphological differentiation with limited clinical usefulness, and on the mucoepidermoid carcinoma, taken as a paradigm of a histotype derived from a preexisting papillary carcinoma through a putative genetic mechanism. It is tempting to suggest that the foci of mucoepidermoid carcinoma that can be found in cases of papillary carcinoma represent a sort of balanced divergent differentiation of the latter. Moreover, we think we have obtained enough evidence to suggest that E-cadherin mutation(s) constitute the genetic basis of such divergent differentiation.

The most-important features of the so-called mixed medullary and follicular (or papillary) carcinoma have been extensively reviewed from a pragmatic standpoint. It is now widely accepted that such mixed carcinomas do, although rarely, exist. However, the controversy surrounding their putative histogenesis(es) is far from settled. Such mixed tumors raise diagnostic problems because, in routine diagnostic work, it is not easy to distinguish a *bona fide* mixed carcinoma from a medullary carcinoma that has absorbed or phagocytosed thyroglobulin. It has also become fashionable to try to subclassify the follicular component of the mixed carcinomas as "true follicular, papillary or poorly differentiated. We have previously stressed that the attempt to characterize the follicle cell derived component of mixed carcinomas contrasts with the usual approach to carcinomas exhibiting divergent differentiation elsewhere in the body. Such an attempt implicitly indicates, on the other hand, that we are not at present sure about the clonal nature of these mixed carcinomas. In other words, the possibility that at least some of them may indeed be collision tumors has to be entertained.

In order to progress in the understanding of the histogenesis and possible molecular mechanisms leading to mixed carcinomas, Volante *et al.* have very recently completed a study of 12 *bona fide* mixed carcinomas of the thyroid, *i.e.*, tumors exhibiting the morphological and immunohistochemical features of both medullary and follicle-cell derived neoplasms. These authors isolated the two histological components of the 12 carcinomas by laser-based microdissection and analyzed them for mutations in the *ret* protooncogene and for allelic losses of nine loci on six chromosomes. In female patients, Volante *et al.* also studied the clonal composition of the carcinomas. The results obtained strongly indicate that the follicular and medullary components in mixed carcinomas are not derived from a single progenitor cell since they consistently exhibited a different pattern of mutations and allelic losses. Furthermore, the seven tumors that were suitable for clonal analysis did not yield any signs of monoclonality. These authors also demonstrated that the follicular structures in mixed carcinomas are often oligo- and polyclonal and more frequently exhibit hyperplastic histological features than neoplastic ones, thus indicating that at least a subset of mixed carcinomas appears to include medullary thyroid carcinomas containing hyperplastic follicles.

It is far beyond the scope of this short presentation to elaborate on the extremely interesting questions raised by the study of Volante *et al.* but we are sure that this seminal work will trigger a new wave of discussion on the mono or dual histogenesis of thyroid tumors.

### References

- Albores-Saavedra H, Gorrses de la Mora T, De la Torre-Rendon F et al. *Mixed medullary-papillary carcinoma of the thyroid: A previously unrecognized variant of thyroid carcinoma.* Hum Pathol 1990; 21: 1151-1155.
- Fonseca E, Sobrinho-Simbes M. *Diagnostic problems in differentiated carcinomas of the thyroid.* Pathol Res Pract 1995; 191: 318-331.
- Li Volsi V. *Surgical pathology of the thyroid.* WB Saunders, Philadelphia 1990.
- Rocha AS, Soares P, Machado JC et al. *Mutations in Mucoepidermoid Carcinoma of the Thyroid (Abstract).* Book of Proceedings of the Meeting on Cell Adhesion and Communication in Growth Control and Cancer. ARC, Lyon 1999.
- Rossi et Rossi J, Csrcangiu M, Oe Lellis R. *Tumors of the Thyroid Gland.* Armed Forces Institute of Pathology, Atlas of Tumor Pathology ~ Series, Washington 1990.
- Sobrinho-Simdes M. *Mixed medullary and follicular carcinoma of the thyroid.* Histopathology 1993; 23: 287-289.
- Sobrinho-Simbes M. *Poody differentiated carcinoma(s) of the thyroid (Editorial).* Endocrine Pathol 1996; 7: 99-102.
- Sobrinho-Simbes M, Nesland JM, Johannessen JV. *Farewell to the dual histogenesis of thyroid tumors.* ultrastruct Pathol 1985; 8(2-3): iii-v.
- Volante M, Papotti M, S peel EJM et al. *Mixed medullary-follicular thyroid carcinomas: Molecular evidence for a dual origin of tumor components.* (Submitted).